

- (ii) conditional density functions ;
- (iii)  $\text{Var}(X)$  ;
- (iv)  $\text{Var}(Y)$  ;
- (v) covariance between  $X$  and  $Y$ .

Total number of printed pages-24

3 (Sem-5/CBCS) MAT HE 1/HE 2/HE 3

2021

(Held in 2022)

**MATHEMATICS**

(Honours Elective)

**Answer the Questions from any one Option.**

**OPTION-A**

Paper : MAT-HE-5016

**( Number Theory )**

**DSE (H)-1**

Full Marks : 80

Time : Three hours

**The figures in the margin indicate full marks for the questions.**

**PART-A**

1. Choose the correct option :  $1 \times 10 = 10$ 
  - (i) Two integers  $a$  and  $b$  are coprime if there exists some integers  $x, y$  such that  
(a)  $ax + by = 1$



- (b)  $ax - by = 1$
- (c)  $(ax + by)^n = 1$
- (d) None of the above

(ii) Let  $d = \gcd(a, b)$ ,  $n \in \mathbb{N}$ . If  $d \mid c$  and  $(x_0, y_0)$  is a solution of linear Diophantine equation  $ax + by = c$ , then all integral solutions are given by

(a)  $(x, y) = \left( x_0 + \frac{bn}{d}, y_0 - \frac{an}{d} \right)$

(b)  $(x, y) = \left( x_0 - \frac{bn}{d}, y_0 + \frac{an}{d} \right)$

(c)  $(x, y) = \left( x_0 + \frac{an}{d}, y_0 - \frac{bn}{d} \right)$

(d)  $(x, y) = \left( x_0 - \frac{an}{d}, y_0 + \frac{bn}{d} \right)$

(iii) A reduced residue system modulo  $m$  is a set of integers  $r_i$  such that

(a)  $[r_i, m] = 1$

(b)  $(r_i, m) = 1$

(c)  $(r_i, m) \neq 1$

(d) None of the above

(iv) Suppose that  $m_j$  are pairwise relatively prime and  $a_j$  are arbitrary integers ( $j = 1, 2, \dots, k$ ) then there exist solution  $x$  to the simultaneous congruence  $x \equiv a_j \pmod{m_j}$ , such that  $x$  are

(a) congruent modulo

$$M = m_1 \cdot m_2 \cdot m_3 \dots m_k$$

(b) congruent modulo  $M = \sum_{j=1}^k m_j$

(c) congruent modulo  $m_i$

(d) Both (a) and (b)

(v) The product of four consecutive positive integers is divisible by

(a) 20

(b) 22

(c) 24

(d) 26

(vi) Euler's  $\phi$ -function of a prime number  $p$ , i.e.,  $\phi(p)$  is

(a)  $p$

(b)  $p - 1$

(c)  $\frac{p}{2} - 1$

(d) None of the above



(vii) For which value of  $m$ ,  
 $\text{CRS}(\text{mod } m) = \text{RRS}(\text{mod } m)$  ?

- (a) If  $m$  is a prime
- (b) If  $m$  is a composite
- (c) If  $m < 10$
- (d) None of the above

(viii) If  $ca \equiv cb(\text{mod } m)$ , then

- (a)  $a \equiv b \left( \text{mod } \frac{m}{(c, m)} \right)$
- (b)  $a \equiv b(\text{mod } m)$
- (c)  $a \equiv b(\text{mod } m, (c, m))$
- (d) None of the above

(ix) The unit place digit of  $2^{73}$  is

- (a) 4
- (b) 6
- (c) 8
- (d) 2

(x) The highest power of 7 that divides  $50!$  is

- (a) 7
- (b) 8
- (c) 10
- (d) 5

2. Answer the following questions :

$$2 \times 5 = 10$$

(a) If  $p$  is a prime, then prove that

$$\phi(p!) = (p-1) \phi((p-1)!) \quad 2$$

(b) Find all prime number  $p$  such that

$$p^2 + 2 \text{ is also a prime.} \quad 2$$

(c) For  $n = p^k$ ,  $p$  is a prime, prove that

$$n = \sum_{d|n} \phi(d)$$

where  $\sum_{d|n}$  denotes the sum over all

positive divisors of  $n$ . 2

(d) Find the number of zeros at the end of the product of first 100 natural numbers. 2

(e) Find  $\sigma(12)$ . 2

3. Answer **any four** questions : 5 \times 4 = 20

(a) If  $\phi$  is Euler's phi function, then find

$$\phi(\phi(1001)). \quad 5$$



(b) Find the remainder, when  $30^{40}$  is divided by 17. 5

(c) State and prove Chinese Remainder Theorem. 5

(d) If  $p_n$  is the  $n$ th prime number, then prove that

$$p_n < 2^{2^{n-1}} \quad 5$$

(e) If  $n = p_1^{k_1} p_2^{k_2} p_3^{k_3} \dots p_r^{k_r}$  is the prime factorization of  $n > 1$ , then prove that

$$(i) \quad \tau(n) = (k_1 + 1)(k_2 + 1)(k_3 + 1) \dots (k_r + 1)$$

$$(ii) \quad \sigma(n) = \frac{p_1^{k_1+1} - 1}{p_1 - 1} \times \frac{p_2^{k_2+1} - 1}{p_2 - 1} \times \dots \times \frac{p_r^{k_r+1} - 1}{p_r - 1}$$

$2\frac{1}{2} + 2\frac{1}{2} = 5$

(f) Define Mobius function. Also show that

$$\mu(m \cdot n) = \mu(m) \cdot \mu(n)$$

Hence find  $\mu(6)$ .  $1+3+1=5$

### PART-B

Answer **any four** questions :  $10 \times 4 = 40$

4. (a) If  $d = (a, n)$ , prove that the linear congruence  $ax \equiv b \pmod{n}$  has a solution if and only if  $d \mid b$ . 5

(b) (i) When a number  $n$  is divided by 3 it leaves remainder 2. Find the remainder when  $3n + 6$  is divided by 3. 2

(ii) Prove that  $5n + 3$  and  $7n + 4$  are coprime to each other for any natural number  $n$ . 3

5. (a) If  $p$  is a prime, then prove that

$$(p-1)! \equiv -1 \pmod{p} \quad 5$$

(b) Using property of congruence show that 41 divides  $2^{20} - 1$ . 5

6. (a) Prove that every positive integer ( $n > 1$ ) can be expressed uniquely as a product of primes. 5

(b) Determine all solutions in the integers of the Diophantine equation  $172x + 20y = 1000$  5

7. (a) If  $n$  be any positive integer and can be expressed as  $n = p_1^{\alpha_1} \cdot p_2^{\alpha_2} \dots p_k^{\alpha_k}$ , then

$$\text{prove that } \phi(n) = n \prod_{j=1}^k \left(1 - \frac{1}{p_j}\right). \quad 5$$



- (b) If  $m$  and  $n$  are any two integers such that  $(m, n) = 1$ , prove that

$$\phi(m.n) = \phi(m). \phi(n). \quad 5$$

8. (a) For each positive integer  $n \geq 1$ , show that

$$\sum_{d|n} \mu(d) = \begin{cases} 1, & \text{if } n = 1 \\ 0, & \text{if } n > 1 \end{cases} \quad 5$$

- (b) If  $k$  denotes the number of distinct prime factors of positive integer  $n$ , then prove that

$$\sum_{d|n} |\mu(d)| = 2^k \quad 5$$

9. (a) Show that  $\sum_{d|n} \mu(d) \tau(d) = (-1)^k$

where  $k$  denotes the number of distinct prime factors of positive integers  $n$ .

5

- (b) Prove that

- (i)  $\tau(n)$  is an odd integer iff  $n$  is a perfect square. 3

- (ii) For any integer  $n \geq 3$ , show that

$$\sum_{k=1}^n \mu(k!) = 1. \quad 2$$

10. (a) Let  $p$  be an odd prime. Show that the congruence  $x^2 \equiv -1 \pmod{p}$  has a solution if and only if  $p \equiv 1 \pmod{4}$ .

5

- (b) If  $n \geq 1$  and  $\gcd(a, n) = 1$ , then prove that  $a^{\phi(n)} \equiv 1 \pmod{n}$ . 5

11. (a) If  $n$  is a positive integer and  $p$  is a prime, then prove that the exponent of the highest power of  $p$  that divides  $n!$

$$\text{is } \sum_{k=1}^{\infty} \left[ \frac{n}{p^k} \right]. \quad 5$$

- (b) Solve  $3[x] = x + 2\{x\}$  where  $[x]$  denotes greatest integer  $\leq x$  and  $\{x\}$  denotes the fractional part of  $x$ . 5



### OPTION-B

Paper : MAT-HE-5026

**(Mechanics)**

Full Marks : 80

Time : Three hours

**The figures in the margin indicate full marks for the questions.**

1. Answer the following questions :  $1 \times 10 = 10$

- (i) What is the physical significance of the moment of a force?
- (ii) State Newton's second law of motion.
- (iii) Define angle of friction.
- (iv) Define the centre of gravity of a body.
- (v) What do you mean by terminal velocity?
- (vi) What is the geometrical representation of the simple harmonic motion?
- (vii) What is the length of arm of a couple equivalent to the couple  $(P, p)$  having constituent force of magnitude  $F$ ?

(viii) Can a force and a couple in the same plane be equivalent to a single force?

(ix) Define a couple.

(x) State Hooke's law.

2. Answer the following questions :  $2 \times 5 = 10$

- (a) Find the greatest and least resultant of two forces acting at a point whose magnitudes are  $P$  and  $Q$  respectively.
- (b) Find the centre of gravity of an arc of a plane curve  $y = f(x)$ .
- (c) State the laws of static friction.
- (d) Show that impulse of a force is equal to the momentum generated by the force in the given time.
- (e) Write the expression for the component of velocity and acceleration along radial and cross-radial direction for a motion of a particle in a plane curve.



3. Answer **any four** questions of the following :  
5×4=20

(a) The line of action of a force  $F$  divides the angle between its component forces  $P$  and  $Q$  in the ratio 1 : 2. Prove that  $Q(F + Q) = P^2$ .

(b)  $P$  and  $Q$  are two like parallel forces. If  $P$  is moved parallel to itself through a distance  $x$ , show that the resultant of  $P$  and  $Q$  moves through a distance  $\frac{Px}{P + Q}$ .

(c)  $R$  is the resultant of two forces  $P$  and  $Q$  acting at a point and at a given angle. If the force  $P$  be doubled, show that the new resultant will be of magnitude  $\sqrt{2(P^2 + R^2) - Q^2}$ .

(d) A particle of mass  $m$  moves in a straight line under acceleration  $mn^2x$  towards a point  $O$  on the line, where  $x$  is the distance from  $O$ . Show that if  $x = a$  and  $\frac{dx}{dt} = u$  when  $t = 0$ , then at time  $t$ ,  
$$x = a \cos nt + \frac{u}{n} \sin nt.$$

(e) A particle moving with simple harmonic motion in a straight line has velocity  $v_1$  and  $v_2$  at distance  $x_1, x_2$  from the centre of its path. Show that if  $T$  be the period

$$\text{of its motion then } T = 2\pi \sqrt{\frac{x_1^2 - x_2^2}{v_2^2 - v_1^2}}.$$

(f) Show that the sum of the kinetic energy and potential energy is constant throughout the motion when a particle of mass  $m$  falls from rest at a height  $h$  above ground.

4. Answer **any four** questions of the following :  
10×4=40

(a) Forces  $P, Q$  and  $R$  act along the sides  $BC, CA$  and  $AB$  of a triangle  $ABC$  and forces  $P', Q'$  and  $R'$  act along  $OA, OB$  and  $OC$ , where  $O$  is the centre of the circumscribed circle, prove that

$$(i) \quad P \cos A + Q \cos B + R \cos C = 0$$

$$(ii) \quad \frac{PP'}{a} + \frac{QQ'}{b} + \frac{RR'}{c} = 0$$



- (b) State and prove Lami's theorem. Forces  $P$ ,  $Q$  and  $R$  acting along  $OA$ ,  $OB$  and  $OC$ , where  $O$  is the circumcentre of triangle  $ABC$ , are in equilibrium. Show that

$$\frac{P}{a^2(b^2 + c^2 - a^2)} = \frac{Q}{b^2(c^2 + a^2 - b^2)} = \frac{R}{c^2(a^2 + b^2 - c^2)}$$

- (c) (i) Find the centre of gravity of a circular arc of radius  $a$  which subtends an angle  $2\alpha$  at the centre.
- (ii) Find the centre of gravity of a uniform parabolic area cut off by a double ordinate at a distance  $h$  from the vertex.
- (d) (i) Show that the least force which will move a weight  $W$  along a rough horizontal plane is  $W \sin \phi$ , where  $\phi$  is the angle of friction.
- (ii) If a body is placed upon a rough inclined plane, and is on the point of sliding down the plane under the action of its weight and the reactions of the plane only, show that the angle of inclination of the plane to the horizon is equal to the angle of friction.

- (e) A particle moves in a straight line under an attraction towards a fixed point on the line varying inversely as the square of the distance from the fixed point. Investigate the motion.
- (f) A particle moves in a straight line  $OA$  starting from the rest at  $A$  and moving with an acceleration which is directed towards  $O$  and varies as the distance from  $O$ . Discuss the motion of the particle. Hence define simple harmonic motion and time period of the motion.
- (g) Find the component of acceleration of a point moving in a plane curve along the initial line and the radius vector. Also find the component of acceleration perpendicular to initial line and perpendicular to radius vector.
- (h) A particle is falling under gravity in a medium whose resistance varies as the velocity. Find the distance and velocity at any time  $t$ . Also find the terminal velocity of the particle.



### OPTION-C

Paper : MAT-HE-5036

#### (Probability and Statistics)

Full Marks : 80

Time : Three hours

**The figures in the margin indicate full marks for the questions.**

1. Answer the following questions :  $1 \times 10 = 10$

(a) Write the sample space for the experiment of tossing a coin three times in succession or tossing three coins at a time.

(b) Is the probability mass function

$x$	-1	0	1
$P(x)$	0.3	0.4	0.4

admissible? Give reason.

(c) Sketch the area under any probability curve with probability function  $p(x)$  between  $x = c$  and  $x = d$  represented by

$$P(c \leq X \leq d) = \int_c^d p(x) dx.$$

(d) What conclusion one can make about the conditional probability  $P(A/B)$  if  $P(B) = 0$ ?

(e) State the multiplicative theorem of expectation.

(f) Mention the relationship among the mean, median and mode of the normal distribution.

(g) If  $X$  and  $Y$  are two independent random variables, then find  $\text{Var}(2X + 3Y)$ .

(h) Write the mean and variance of standard normal variate  $Z = \frac{X - \mu}{\sigma}$ ,

where  $\mu$  and  $\sigma$  are mean and standard deviation respectively.

(i) When is the correlation coefficient between two random variables  $X$  and  $Y$  zero.

(j) State weak law of large number.



2. Answer the following questions :  $2 \times 5 = 10$

- (a) Prove that probability of any impossible event is zero.
- (b) If  $X$  is a random variable, then prove that  $\text{Var}(X) = E(X^2) - \{E(X)\}^2$
- (c) Find the constant  $c$  such that the function

$$f(x) = cx^2, 0 < x < 3 \\ 0, \text{ otherwise}$$

is a density function and also find  $P(0 < x < 3)$ .

- (d) A random variable  $X$  has density function given by

$$f(x) = 2e^{-2x}, x \geq 0 \\ 0, \text{ otherwise}$$

then find the moment generating function.

- (e) Comment on the following statement :  
"The mean of a binomial distribution is 3 and its standard deviation is 2".

3. Answer **any four** parts from the following :  
 $5 \times 4 = 20$

- (a) A bag contains 6 white and 9 black balls. Four balls are drawn at a time. Find the probability for the first draw to give 4 white and the second to give 4 black balls if the balls are not replaced before the second draw.

- (b) For two independent events  $A$  and  $B$  prove that (i)  $A$  and  $B$  are independent, and (ii)  $\bar{A}$  and  $\bar{B}$  are independent.

- (c) A random variable  $X$  has the function

$$f(x) = \frac{c}{x^2 + 1}, \text{ where } -\infty < x < \infty, \text{ then}$$

- (i) find the value of the constant  $c$  ;  
(ii) find the probability that  $X^2$  lies between  $\frac{1}{3}$  and 1.

- (d) The joint probability of two variables  $X$  and  $Y$  is given by

$$f(x, y) = \frac{1}{42} (2x + y), 0 \leq x \leq 2, 0 \leq y \leq 3 \\ 0, \text{ otherwise}$$

Find (i)  $F(Y/2)$ , and (ii)  $P(y = 1/x = 3)$



(e) A coin is tossed until a head appears. What is the expectation of the number of tosses required?

(f) The probability of a man hitting a target is  $\frac{1}{4}$ .

(i) If he fires 7 times, what is the probability of his hitting the target at least twice?

(ii) How many times must he fire so that the probability of his hitting the target at least once is greater than  $\frac{2}{3}$ ?

4. Answer **any four** parts from the following:  
 $10 \times 4 = 40$

(a) For  $n$  events  $A_1, A_2, A_3, \dots, A_n$ , prove that

$$P\left(\bigcup_{i=1}^n A_i\right) = \sum_{i=1}^n P(A_i) - \sum_{1 \leq i < j \leq n} P(A_i \cap A_j) + \sum_{1 \leq i < j < k \leq n} P(A_i \cap A_j \cap A_k) + \dots + (-1)^{n-1} P(A_1 \cap A_2 \cap A_3 \dots \cap A_n)$$

Hence find  $P\left(\bigcup_{i=1}^3 A_i\right)$

(b) Suppose that two dimensional continuous random variables  $(X, Y)$  has joint p.d.f given by

$$f(x, y) = 6x^2y, \quad 0 < x < 1, \quad 0 < y < 1 \\ 0, \quad \text{elsewhere}$$

(i) Verify that  $\int_0^1 \int_0^1 f(x, y) dx dy = 1$

(ii) Find  $P\left(0 < X < \frac{3}{4}, \frac{1}{3} < Y < 2\right)$

(iii) Find  $P(X + Y < 1)$

(iv) Find  $P(X > Y)$

(v) Find  $P(X < 1/Y < 2)$

(c) (i) The probability function of a random variable  $X$  is given by

$$f(x, y) = \frac{x^2}{81}, \quad -3 < x < 6 \\ 0, \quad \text{otherwise}$$

Find the probability density function for the random variable

$$u = \frac{1}{3}(12 - X).$$



- (ii) Define moment generating function of a random variable  $X$ . Find the moment generating function of binomial distribution.

- (d) (i) The probability curve  $y = f(x)$  has a range from 0 to  $\infty$ . If  $f(x) = e^{-x}$ , find the mean and variance.

- (ii) If  $X$  be a continuous random variable with probability density function

$$f(x) = ax, 0 \leq x \leq 1$$

$$a, 1 \leq x \leq 2$$

$$-ax + 3a, 2 \leq x \leq 3$$

$$0, \text{ otherwise}$$

compute  $P(X \leq 1.5)$

- (e) (i) Derive Poisson distribution as a limiting case of binomial distribution.

- (ii) Prove that mean and variance of a binomially distributed variable are respectively  $np$  and  $npq$ .

- (f) (i) Define correlation coefficient of two random variables  $X$  and  $Y$ . Show that correlation coefficient is independent of change of origin and scale.

- (ii) Obtain the equation of two lines of regression for the following data :

$X :$	65	66	67	67	68	69	70	72
$Y :$	67	68	65	68	72	72	69	71

Also obtain the estimate of  $X$  for  $Y = 70$ .

- (g) (i) If  $X$  is a random variable with mean  $\mu$  and variance  $\sigma^2$ , then for any positive number  $k$ , prove that

$$P\{|X - \mu| \geq k\sigma\} \leq \frac{1}{k^2}$$

- (ii) A symmetric die is thrown 600 times. Find the lower bound for the probability of getting 80 to 120 sixes.

- (h) The random variables  $X$  and  $Y$  have the following joint probability density function :

$$f(x, y) = 2 - x - y; 0 \leq x \leq 1, 0 \leq y \leq 1$$

$$0, \text{ otherwise}$$

Find—

- (i) marginal probability density functions of  $X$  and  $Y$ ;



**Or**

Explain with example the 'if' statement and nested 'if' statement in C. Write a C program to find the roots of a quadratic equation  $ax^2 + bx + c = 0$ , for all possible values of  $a, b, c$ . 5+5=10

6. What is the basic difference between 'Library functions' and 'User-defined functions'? Mention *two* advantages of using 'User-defined functions'. How are such functions declared and called in a program? Write a C program using function to find the biggest of three numbers. 1+2+2+5=10

**Or**

Write a C programme that reads a number, obtains a new number by reversing the digits of the given number, and then determine the gcd of the two numbers. To build the programme, use two functions — one to find gcd and another to reverse the digits. 10

Total number of printed pages-16

**3 (Sem-5/CBCS) MAT HE 4/5/6**

**2021**

**(Held in 2022)**

**MATHEMATICS**

**(Honours Elective)**

**Answer the Questions from any one Option.**

**OPTION-A**

Paper : MAT-HE-5046

**(Linear Programming)**

**DSE(H)-2**

Full Marks : 80

Time : Three hours

***The figures in the margin indicate full marks for the questions.***

1. Answer the following as directed : 1×10=10
- (a) A basic feasible solution whose variables are.
- (i) degenerate
  - (ii) nondegenerate

*Contd.*



(iii) non-negative

(iv) None of the above

*(Choose the correct answer)*

(b) The inequality constraints of an LPP can be converted into equation by introducing

(i) negative variables

(ii) non-degenerate B.F.

(iii) slack and surplus variables

(iv) None of the above

*(Choose the correct answer)*

(c) A solution of an LPP, which optimize the objective function is called

(i) basic solution

(ii) basic feasible solution

(iii) optimal solution

(iv) None of the above

*(Choose the correct answer)*

(d) What is artificial variable of an LPP ?

(e) Write the equation of line segment in  $\mathbb{R}^n$ .

(f) Define dual of a given LPP.

(g) What is pure strategy of game theory ?

(h) Is region of feasible solution to an LPP constitute a convex set ?

(i) Is every convex set in  $\mathbb{R}^n$  a convex polyhedron also ?

(j) Is every boundary point an extreme point of a convex set ?

2. Answer the following questions :  $2 \times 5 = 10$

(a) Show that the feasible solution

$x_1 = 1, x_2 = 0, x_3 = 1, z = 6$  to the system

$$\min Z = 2x_1 + 3x_2 + 4x_3$$

$$\text{s.t. } x_1 + x_2 + x_3 = 2$$

$$x_1 - x_2 + x_3 = 2, \quad x_i \geq 0$$

is not basic.

(b) A hyperplane is given by the equation

$$3x_1 + 2x_2 + 4x_3 + 7x_4 = 8$$

Find in which half space do the point  $(-6, 1, 7, 2)$  lie.

(c) Find extreme points if any of the set  $S = \{(x, y) : |x| \leq 1, |y| \leq 1\}$

(d) Show by an example that the union of two convex sets is not necessarily a convex set.



(e) If  $x_1 = 2, x_2 = 3, x_3 = 1$  a BFS of the LPP

$$\max Z = x_1 + 2x_2 + 4x_3$$

$$\text{s.t. } 2x_1 + x_2 + 4x_3 = 11$$

$$3x_1 + x_2 + 5x_3 = 14$$

$$x_1, x_2, x_3 \geq 0 ? \text{ Explain.}$$

3. Answer **any four** questions :  $5 \times 4 = 20$

(a) Prove that the set of all feasible solutions of an LPP is a convex set.

(b) Sketch the convex polygon spanned by the following points in a two-dimensional Euclidean space. Which of these points are vertices ? Express the other as the convex linear combination of the vertices

$$(0,0), (0,1), (1,0), \left(\frac{1}{2}, \frac{1}{4}\right).$$

(c) If  $x_0 \in S$  where  $S$  is the set of all FS of the LPP  $\min Z = cx$ , such that  $Ax = b, x \geq 0$  minimize the objective function  $Z = cx$ , then show that  $x_0$  also maximize the objective function  $Z^* = (-c)x$  over  $S$ .

(d) Find the dual of the following LPP :

$$\min Z_p = x_1 + x_2 + x_3$$

$$\text{s.t. } x_1 - 3x_2 + 4x_3 = 5$$

$$2x_1 - 3x_2 \leq 3$$

$$2x_2 - x_3 \geq 5$$

$$x_1, x_2, x_3 \geq 0$$

(e) Prove that the dual of a dual is a primal problem itself.

(f) Write the characteristics of an LPP in canonical form.

4. Answer (a) **or** (b), (c) **or** (d), (e) **or** (f),  
(g) **or** (h) :  $10 \times 4 = 40$

(a) Old hens can be bought for Rs. 2 each but young ones cost Rs. 5 each. The old hens lay 3 eggs per week and the young ones 5 eggs per week, each being worth 30 paise. A hen costs Re. 1 per week to feed. If I have only Rs. 80 to spend for hens, how many of each kind shall I buy to give a profit of more than Rs. 6 per week, assuming that I can not house more than 20 hens ? Formulate the LPP and solve by graphical method.

(b) Find all basic and then all the basic feasible solutions for the equations

$$2x_1 + 6x_2 + 2x_3 + x_4 = 3$$

$$6x_1 + 4x_2 + 4x_3 + 6x_4 = 2$$

and determine the associated general convex combination of the extreme point solutions.



(c) State and prove the fundamental theorem of LPP.

(d) Solve by simplex method :

$$\max Z = 3x_1 + 5x_2 + 4x_3$$

$$\text{s.t. } 2x_1 + 3x_2 \leq 8$$

$$3x_1 + 2x_2 + 4x_3 \leq 15$$

$$2x_2 + 5x_3 \leq 10$$

$$x_1, x_2, x_3 \geq 0$$

(e) If in an assignment problem, a constant is added or subtracted to every element of a row (or column) of the cost matrix  $[c_{ij}]$ , then prove that an assignment which minimizes the total cost for one matrix, also minimizes the total cost for the other matrix.

(f) Solve the following transportation problem :

		To				
		$S_1$	$S_2$	$S_3$	$S_4$	Supply
From	$O_1$	1	2	1	4	30
	$O_2$	3	3	2	1	50
	$O_3$	4	2	5	9	20
Demand		20	40	30	10	100

(g) For any zero-sum two-persons game where the optimal strategies are not pure and for which A's pay-off matrix is

		B	
		$Iy_1$	$IIy_2$
A	$x_1 I$	$a_{11}$	$a_{12}$
	$x_2 II$	$a_{21}$	$a_{22}$

the optimal strategies are  $(x_1, x_2)$  and  $(y_1, y_2)$  then prove that

$$\frac{x_1}{x_2} = \frac{a_{22} - a_{21}}{a_{11} - a_{12}} \quad \text{and} \quad \frac{y_1}{y_2} = \frac{a_{22} - a_{12}}{a_{11} - a_{21}} \quad \text{and}$$

the value of the game to A is given by

$$v = \frac{a_{11}a_{22} - a_{12}a_{21}}{(a_{11} + a_{22}) - (a_{12} + a_{21})}$$

(h) Solve the game whose pay-off matrix is

-1	-2	8
7	5	-1
6	0	12



## OPTION-B

Paper : MAT-HE-5056

### ( Spherical Trigonometry and Astronomy )

Full Marks : 80

Time : Three hours

**The figures in the margin indicate full marks for the questions.**

1. Answer the following questions :  $1 \times 10 = 10$

- (a) State *one* fundamental difference between a spherical triangle and a plane triangle.
- (b) Define primary circle.
- (c) Define polar triangle and its primitive triangle.
- (d) State the third law of Kepler.
- (e) Explain what is meant by rising and setting of stars.
- (f) Write *any two* coordinate systems to locate the position of a heavenly body on the celestial sphere.
- (g) Define synodic period of a planet.
- (h) Mention *one* property of pole of a great circle.

- (i) Just mention how a spherical triangle is formed.
- (j) What is the declination of the pole of the ecliptic ?

2. Answer the following questions :  $2 \times 5 = 10$

- (a) Prove that section of a sphere by a plane is a circle.
- (b) Discuss the effect of refraction on sunrise.
- (c) Drawing a neat diagram, discuss how horizontal coordinates of a heavenly body are measured.
- (d) Prove that the altitude of the celestial pole at any place is equal to the latitude of that place.
- (e) Show that right ascension  $\alpha$  and declination  $\delta$  of the sun is always connected by the equation  $\tan \delta = \tan \varepsilon \sin \alpha$ ,  $\varepsilon$  being obliquity of the ecliptic.

3. Answer **any four** of the following :

$5 \times 4 = 20$

- (a) Deduce Kepler's laws from Newton's law of gravitation.



(b) Show that the velocity of a planet in its elliptic orbit is  $v^2 = \mu \left( \frac{2}{r} - \frac{1}{a} \right)$  where

$\mu = G(M + m)$  and  $a$  is the semi-major axis of the orbit.

(c) If  $z_1$  and  $z_2$  are the zenith distances of a star on the meridian and the prime vertical respectively, prove that

$$\cot \delta = \operatorname{cosec} z_1 \sec z_2 - \cos z_1$$

where  $\delta$  is the star's declination.

(d) If  $H$  be the hour angle of a star of declination  $\delta$  when its azimuth is  $A$  and  $H'$  when the azimuth is  $(180^\circ + A)$ , show that

$$\tan \phi = \frac{\cos \frac{1}{2}(H' + H)}{\cos \frac{1}{2}(H' - H)}$$

(e) In an equilateral spherical triangle  $ABC$ , prove that  $2 \cos \frac{a}{2} \sin \frac{A}{2} = 1$ .

(f) If  $\psi$  is the angle which a star makes at rising with the horizon, prove that  $\cos \psi = \sin \phi \sec \delta$ , where the symbols have their usual meanings.

4. Answer **any four** questions of the following : 10×4=40

(a) If the colatitude is  $C$ , prove that

$$C = x + \cos^{-1}(\cos x \sec y)$$

where  $\tan x = \cot \delta \cos H$  and

$$\sin y = \cos \delta \sin H,$$

$H$  being the hour angle.

(b) In any spherical triangle  $ABC$ , prove that  $\frac{\sin A}{\sin a} = \frac{\sin B}{\sin b} = \frac{\sin C}{\sin c}$ . Also prove

$$\text{that } \frac{\sin(A+B)}{\sin C} = \frac{\cos a + \cos b}{1 + \cos c}$$

(c) Define astronomical refraction and state the laws of refraction. Derive the formula for refraction as  $R = k \tan \xi$ ,  $\xi$  being the apparent zenith distance of a heavenly body. Mention *one* limitation of this formula.

(d) On account of refraction, the circular disc of the sun appears to be an ellipse. Prove it.

(e) Derive Kepler's equation in the form  $M = E - e \sin E$ , where  $M$  and  $E$  are respectively mean anomaly and eccentric anomaly.



- (f) Assuming the planetary orbits to be circular and coplanar, prove that the sidereal period  $P$  and the synodic period  $S$  of an inferior planet are related to the earth's periodic time  $E$  by

$$\frac{1}{S} = \frac{1}{P} - \frac{1}{E}$$

Calculate the sidereal period (in mean solar days) of a planet whose sidereal period is same as its synodic period.

- (g) Prove that, if the fourth and higher powers of  $e$  are neglected,

$$E = M + \frac{e \sin M}{1 - e \cos M} - \frac{1}{2} \left( \frac{e \sin M}{1 - e \cos M} \right)^3$$

is a solution of Kepler's equation in the form.

- (h) Derive the expressions to show the effect of refraction in right ascension and declination.

## OPTION-C

Paper : MAT-HE-5066

( *Programming in C* )

Full Marks : 60

Time : Three hours

**The figures in the margin indicate full marks for the questions.**

1. Answer the following questions : 1×7=7

- (a) Write *any two* special characters that are used in C.
- (b) Mention *two* data types that are used in C language.
- (c) For  $x = 2$ ,  $y = 5$ , write the output of the C function 'pow ( $x$ ,  $y$ )'.
- (d) Convert the mathematical expression 
$$z = e^x + \log y + \sqrt{1+x}$$
 into C expression.
- (e) Write the utility of clrscr ( ) function.
- (f) Write a difference between local variable and global variable.
- (g) Write the C library function which can evaluate  $|x|$ .



2. Answer the following questions :  $2 \times 4 = 8$

- (a) Write the difference between 'assignment' and 'equality'.
- (b) How does ' $x + +$ ' differ from ' $+ + x$ ' ?
- (c) What is a string constant ? Give an example.
- (d) Write *four* relational operators that are used in C.

3. Answer **any three** parts :  $5 \times 3 = 15$

- (a) Explain arithmetic and logical operators in C with suitable examples.
- (b) List three header files that are used in C. Also write their utilities.  $3 + 2 = 5$

$A = 5; B = 3$

$A = A + B;$

$B = A - B;$

$A = A - B;$

Write the output of A and B from the above program segment in C.

- (c) Write a C program to find the sum of all odd integers between 1 and  $n$ .
- (d) Write the general form of do-while loop and explain how it works with the help of a suitable example.

- (e) Write the utility of 'break' and 'continue' statements with the help of suitable examples.

4. Why are arrays required in C programming ? How are one-dimensional arrays declared and inputs given to array ? Explain briefly with example. Write a program to read given  $n$  numbers and then find the sum of all positive and negative numbers.  $2 + 3 + 5 = 10$

**Or**

How are two-dimensional arrays declared ? Write a C program to read a  $3 \times 3$  matrix and print the same as output. Hence write a C program to read a  $3 \times 3$  matrix, print its transpose and write the determinants of both.  $1 + 4 + 5 = 10$

5. Write a C program for each of the following :

- (a) To evaluate the function  $5$   
 $f(x) = x^2 + 2x - 10, x \geq 0$   
 $= |x|, x < 0$

- (b) To find the biggest of three numbers.  $5$